

Phases of Wave Functions and Level Repulsion

W.D. Heiss

Centre for Nonlinear Studies and Department of Physics
University of the Witwatersrand, PO Wits 2050, Johannesburg, South Africa

PACS: 03.65.Bz, 02.30.Dk, 84.40.-x

Avoided level crossings are associated with exceptional points which are the singularities of the spectrum and eigenfunctions, when they are considered as functions of a coupling parameter. It is shown that the wave function of *one* state changes sign but not the other, if the exceptional point is encircled in the complex plane. An experimental setup is suggested where this peculiar phase change could be observed.

The phase of the wave function changes in a characteristic way, if a selfadjoint Hamiltonian has an energy degeneracy at specific values in some parameter space and if a loop around the point of degeneracy is described in parameter space [1]. In the simple case of a real symmetric Hamiltonian two parameters are needed to get a *diaboloic point*. In this case Berry's phase is π when looping around the degeneracy in two dimensional parameter space.

If a selfadjoint Hamiltonian depends only on one parameter, its variation will in general give rise to level repulsion [2]. Associated with a level repulsion is a pair of exceptional points [3] which are the points where the two levels actually coalesce when continued analytically into the complex plane of the parameter [4]. In the present letter we discuss the behaviour of the wave functions and their phases when an exceptional point is encircled. We suggest an experimental situation where such behaviour could be measured. It is distinctly different from a diaboloic point where Berry's phase occurs. In particular, an exceptional point must not be confused with a coincidental degeneracy of resonance states, which was considered in [5] as a generalisation of Berry's phase.

The difference between a diaboloic point and an exceptional point is due to the selfadjointness of the Hamiltonian in the former and the lack of it in the latter case. Also, when continuing into the complex parameter plane we are faced with analytic functions of a complex variable which implies a more rigid mathematical structure. In the references quoted above a thorough discussion is given of the spectrum and eigenfunctions, when one parameter on which the Hamiltonian depends is continued into the complex plane. The parameter chosen can be an interaction strength as considered in this paper, but other choices are possible [6]. Generically, an N -dimensional matrix problem yields $N(N-1)$ exceptional points. For an infinite dimensional problem, an infinite number can

occur [7], but, depending on the particular situation, they may accumulate at one point in the finite plane [8]; there are special cases, where no exceptional point occurs in the finite plane as can be seen from the analytic solution for the single particle motion in the Hulthen potential [9].




FIG. 1. Level repulsion using in Eq.(3) $\epsilon_1 = 1$, $\epsilon_2 = 2$, $\omega_1 = 2$, $\omega_2 = -1$ and $\phi = \pi/25$.

All essential aspects of exceptional points can be illustrated in a two level model on an elementary level. In fact, for finite or infinite dimensional problems an isolated exceptional point can be described locally by a two dimensional problem [10]. In other words, even though a high or infinite dimensional problem is globally more complex than the two dimensional problem, we do not lose generality for our specific purpose when the restriction to a two dimensional problem is made. For easy illustration we therefore confine ourselves to the discussion of

$$H = \begin{pmatrix} \epsilon_1 & 0 \\ 0 & \epsilon_2 \end{pmatrix} + \lambda U \begin{pmatrix} \omega_1 & 0 \\ 0 & \omega_2 \end{pmatrix} U^\dagger \quad (1)$$

with

$$U = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}. \quad (2)$$

This is, up to a similarity transformation, the most general form of a real two dimensional Hamilton matrix of the type $H_0 + \lambda H_1$. The particular dependence on the parameter λ has been chosen as it is of a nature widely used in physical applications. We emphasise that our aim is not in particular directed at a physical model that is describable by a two dimensional problem although there may exist interesting problems in our special context. The restriction has been chosen for illustration, while the physical application that we have in mind is an

infinite dimensional situation.




FIG. 2. Contours in the complex energy plane. The top illustrates contours for each level, if a closed loop is described in the λ -plane without encircling the exceptional point. The bottom illustrates the energy contour produced by two (equal) closed loops in the λ -plane, which encircle the exceptional point. The solid line corresponds to the first closed loop in the λ -plane and the dotted line to the subsequent loop. The solid dot is the position where the contours would meet, if the loops would cross the exceptional point in the λ -plane.

The eigenvalues of H are given by

$$E_{1,2}(\lambda) = \frac{\epsilon_1 + \epsilon_2 + \lambda(\omega_1 + \omega_2)}{2} \pm R \quad (3)$$

where

$$R = \left\{ \left(\frac{\epsilon_1 - \epsilon_2}{2} \right)^2 + \left(\frac{\lambda(\omega_1 - \omega_2)}{2} \right)^2 + \frac{1}{2} \lambda(\epsilon_1 - \epsilon_2)(\omega_1 - \omega_2) \cos 2\phi \right\}^{1/2}. \quad (4)$$

Clearly, when $\phi = 0$ the spectrum is given by the two lines

$$E_k^0(\lambda) = \epsilon_k + \lambda\omega_k, \quad k = 1, 2$$

which intersect at the point of degeneracy $\lambda = -(\epsilon_1 - \epsilon_2)/(\omega_1 - \omega_2)$. When the coupling between the two levels is turned on by switching on ϕ , the degeneracy is lifted and avoided level crossing occurs as is illustrated in Fig.1. Now the two levels coalesce in the complex λ -plane where R vanishes which happens at the complex conjugate points

$$\lambda_c = -\frac{\epsilon_1 - \epsilon_2}{\omega_1 - \omega_2} \exp(\pm 2i\phi). \quad (5)$$

At these points, the two levels $E_k(\lambda)$ are connected by a square root branch point, in fact the two levels are the values of one analytic function on two different Riemann sheets. In Fig.2a we display contours in the complex energy plane for each level, which are obtained if a loop is described in the complex λ -plane, which does not encircle the exceptional point. Correspondingly, Fig.2b shows the contours, if the exceptional point is encircled. In this case only a double loop in the λ -plane yields a closed loop in the energy plane. Obviously, this connection is not of the type encountered at a genuine diabolic point.

The difference has a bearing also on the scattering matrix [11] and on the wave functions $\psi_1(\lambda)$ and $\psi_2(\lambda)$. In [11], although the notion *exceptional points* is not used, the pertinent distinction between a genuine degeneracy of two resonances and the analytic coalescence (exceptional point) of two (complex) eigenvalues is nicely discussed. A usual degeneracy of two resonances still gives rise to a simple pole in the scattering matrix or Green's function, while an exceptional point produces a double pole. With regard to the eigenfunctions, we recall that for complex λ the Hamiltonian is no longer selfadjoint. This means that the eigenfunctions are no longer orthogonal. We rather have a biorthogonal system which can be normalized as

$$\langle \tilde{\psi}_1(\lambda) | \psi_2(\lambda) \rangle = \delta_{1,2} \quad (6)$$

where $|\psi_k\rangle$ and $\langle \tilde{\psi}_k|$ are the right hand and left hand eigenvectors of H , respectively. Note that Eq.(6) causes problems at the exceptional point, since it is exactly at this point where two linearly independent eigenfunctions no longer exist. This is in contrast to a genuine degeneracy of a selfadjoint operator where a k -fold degeneracy always gives rise to a k -dimensional eigenspace. At the exceptional point, not only the eigenvalues but also the eigenfunctions coalesce. As a consequence, the orthogonality conflicts with the normalization, in other words, if Eq.(6) is enforced globally, that is also at $\lambda = \lambda_c$, the components of the wave function have to blow up. This can be made explicit by parametrizing the wave functions by the complex angle θ , viz.

$$\psi_1(\lambda) = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \quad \psi_2(\lambda) = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (7)$$

with

$$\tan^2 \theta(\lambda) = \frac{E_1(\lambda) - E_2(\lambda) - (\epsilon_1 - \epsilon_2) - \lambda(\omega_1 - \omega_2) \cos 2\phi}{E_1(\lambda) - E_2(\lambda) + (\epsilon_1 - \epsilon_2) + \lambda(\omega_1 - \omega_2) \cos 2\phi}. \quad (8)$$

At $\lambda = \lambda_c$ it is $E_1 = E_2$ and hence $\tan^2 \theta = -1$ which implies $|\cos \theta| = |\sin \theta| = \infty$, that is the components of the wave functions blow up. (Note that $\tan^2 \theta \equiv 0$ in the

trivial case $\phi = 0$). The increase of the components of the wave functions while approaching exceptional points has been used in similar context as a theoretical signature of a phase transition [8], but we do not believe that it has observational consequences.

We now study the behaviour of the wave functions in more detail for two contours in the λ -plane which start, say, at $\lambda = 0$ and end at large real values of λ , but enclose an exceptional point between the two contours. For the complex angle θ we choose an expression which is more convenient for this purpose, *viz.*

$$\tan \theta(\lambda) = \frac{\lambda(\omega_1 - \omega_2) \sin 2\phi}{E_1(\lambda) - E_2(\lambda) + \epsilon_1 - \epsilon_2 + \lambda(\omega_1 - \omega_2) \cos 2\phi}. \quad (9)$$

The first path can be taken along the real λ -axis. From Eq.(9) we read off the expected result that $\theta(0) = 0$ and $\theta(\lambda) \rightarrow \phi$ for $\lambda \gg |(\epsilon_1 - \epsilon_2)/(\omega_1 - \omega_2)|$. In obtaining this result use is made of $E_1 - E_2 = 2R \rightarrow \lambda(\omega_1 - \omega_2)$ for $\lambda \gg |(\epsilon_1 - \epsilon_2)/(\omega_1 - \omega_2)|$. For the second path we move into the upper λ -plane in order to pass above the exceptional point before returning down to the real axis again. Using again Eq.(9) we now have to observe that we crossed into the other sheet which means $E_1 - E_2 = -2R \rightarrow -\lambda(\omega_1 - \omega_2)$. As a consequence we find this time $\tan \theta = -\cot \phi = \tan(\phi + \pi/2)$. Surely we would expect the wave functions to interchange just like the energies, if an exceptional point is encircled. But our finding indicates that *one wave function has changed its sign*. In fact, we obtain along the second path

$$\psi_1 \rightarrow \psi_2, \quad \psi_2 \rightarrow -\psi_1. \quad (10)$$

Note that this result is equally obtained, if a closed contour which encloses the exceptional point is described in the λ -plane. Of interest is the result of a double loop in the λ -plane. It yields

$$\psi_1 \rightarrow -\psi_1, \quad \psi_2 \rightarrow -\psi_2 \quad (11)$$

which is, in accordance with the corresponding single loop in the energy plane, just Berry's phase retrieved. We mention that this implies that the wave functions, if parametrized as in Eq.(7), have an algebraic singularity that is determined by a fourth root at the exceptional point; only a four-fold loop in the λ -plane restores completely the original situation.

Next we address the question concerning the physical significance of these findings. It boils down to the problem of varying a *complex* interaction parameter in the laboratory. Problems of a similar nature have been discussed in connection with Berry's phase for dissipative systems [12], but not as yet for exceptional points. We are guided by the phenomenological description of open quantum systems [8] and submit as one suggestion a strongly absorptive system, where the parameter λ in

Eq.(1) is traditionally replaced by $-iG$ with real absorption parameter G . With the replacement the eigenvalues acquire imaginary parts which are related to the inverse life times of the states of the open system. The exceptional points appear now in the complex G -plane at

$$G_c = -\frac{\epsilon_1 - \epsilon_2}{\omega_1 - \omega_2} \exp(\pm 2i\phi + i\pi/2). \quad (12)$$

If the coupling of the two channels is equal, that is if $\phi = \pi/4$, the two exceptional points lie on the real G -axis at

$$G_c = \pm \frac{\epsilon_1 - \epsilon_2}{\omega_1 - \omega_2}.$$

If the coupling is nearly equal, they lie just above or below the real axis depending on the coupling being slightly weaker or stronger ($\phi < \pi/4$ or $\phi > \pi/4$). Controlling the absorption parameter G and the relative coupling enables one to pass the exceptional point on a path below and above, which is the situation described above. A setup similar to the one proposed by Berry [1] and implemented in [13] should make feasible a distinction of the two cases. In a two dimensional electromagnetic resonator the relative coupling of two suitable levels, that is levels that do coalesce, can be controlled by a judicious choice of the geometry of the resonator [13]. The global absorption can be controlled by radiation losses regulated by suitable antennas. In this way the change of the phase of *one but not the other* wave function should be observable. Note that, for large values of G , one state will be much broader than the other [8].

We stress again that the occurrence of exceptional points is a generic mathematical feature associated with any system that has avoided level crossing. While this presentation used a two dimensional illustration, the results can be immediately generalised. In fact, if more than one exceptional point is encircled, the resulting phase change is simply a combination of several two-dimensional cases. The findings presented in this paper have therefore universal significance. Whether the particular phase changes of the respective wave functions are detectable, is subject to experimentation. Due to the short life time involved for one of the states, scattering experiments, where a steady incoming flux can be maintained, are expected to be particularly suitable. Fortunately, in the spirit of the experiment performed by [13] the adiabatic change of the parameters is not essential, since it is the movement of the nodes of the wave functions that are being observed under parameter variation [14]. Possible jumps of the phases do therefore not affect the result of interest.

- [1] M.V. Berry, *Quantum Chaos*, ed. by G. Casati (London: Plenum) 1985; Proc. R. Soc. A **239**, 45 (1983)
- [2] J. von Neumann and E. Wigner, Z.Phys. **30**, 467 (1929)
- [3] T. Kato, *Perturbation theory of linear operators* Springer, Berlin 1966
- [4] W.D. Heiss and A.L. Sannino, J. Phys. A **23** 1167 (1990); W.D. Heiss and A.L. Sannino, Phys. Rev. A **43** 4159 (1991)
- [5] C. Mimiatura, C. Sire, J. Baudon and J. Bellissard, Eur.Lett. **13**, 199 (1990)
- [6] C. M. Bender and S. Boettcher, Phys.Rev.Lett. **80** 5243 (1998); S.C. Creagh and N.D. Whelan, *chaodyn/9808014*
- [7] W.D. Heiss, Nucl.Phys. **A169** 248 (1971)
- [8] W.D. Heiss, M. Müller and I. Rotter, Phys.Rev. **E**, to appear
- [9] R.G. Newton, *Scattering Theory of Waves and Particles*, Section 14.4, McGraw-Hill 1966
- [10] W.D. Heiss and W.-H. Steeb, J. Math. Phys. **32**, 3003 (1991)
- [11] E. Hernandez and A. Mondragon, Phys.Lett. B **326**, 1, (1994); A. Mondragon and E. Hernandez, J. Phys. A **26**, 5595 (1993)
- [12] G. Nenciu and G. Rasche, Journal of Physics A (Mathematical and General) **25** 5741 (1992); T.B. Kepler and M.L. Kagan, Phys.Rev.Lett. **66** 847 (1991); D. Ellinas, S.M. Barnett and M.A. Dupertuis, Phys.Rev. A **39** 3228 (1989)
- [13] H.-M. Lauber, P. Weidenhammer and D. Dubbers, Phys.Rev.Lett. **72** 1004 (1994)
- [14] H.-D. Gräf, Technische Universität Darmstadt, private communication.